

Healthcare Costs Among Commercially-Insured Patients with Hereditary Angioedema Managed with Long-Term Prophylaxis: A Retrospective US Claims Database Analysis

Raffi Tachdjian¹, Daniel Soteres², Rose Chang³, Manasi Mohan³, Megan Pinaire³, Maral DerSarkissian³, Vibha Desai⁴, Alice Wang⁴, Paul Audhya⁴

¹David Geffen School of Medicine, UCLA, Santa Monica, CA, USA; ²Asthma & Allergy Associates, PC and Research Center, Colorado Springs, CO, USA; ³Analysis Group, Inc., Boston, MA, USA; ⁴KalVista Pharmaceuticals, Inc., Cambridge, MA, USA


Background

- Hereditary angioedema (HAE) is a rare genetic disease associated with unpredictable, painful, and debilitating attacks of tissue swelling in various locations of the body that can be life-threatening depending on the location(s) affected
- Management of HAE consists of lifestyle management, on-demand treatment, and, for appropriate patients, the addition of long-term prophylaxis (LTP)
- Although reductions in HAE attack frequency have been demonstrated with LTP, most patients continue to experience attacks requiring on-demand treatment¹
- Given the increasing number of patients receiving non-androgen LTP and the limited data on related real-world healthcare costs in the US, we estimated these costs using a large retrospective insurance claims database

Methods

- We conducted a retrospective study using the IQVIA PharMetrics® Plus Database (April 1, 2017—March 31, 2022)
- As there are no HAE-specific International Classification of Diseases diagnosis codes, we used a claims-based algorithm to identify potential patients, which required either ≥1 claim for exclusive HAE therapies or ≥2 claims with HAE-related codes with ≥1 claim for other HAE medications²
- Patients were ≥12 years old at index, commercially insured, had ≥6 months of continuous enrollment before and ≥3 months following index
 - Index date = date of first claim for a non-androgen LTP therapy
 - Follow-up period: patients were followed from index date until the earliest of either health plan disenrollment or end of data availability
- This analysis examined outcomes among commercially-insured patients receiving non-androgen LTP (lanadelumab, berotralstat, intravenous [IV] and subcutaneous [SC] C1 inhibitor [C1-INH])
 - Patients may have been treated with more than one LTP during the study period; given the small sample size, the outcomes of each cohort were described based on the index treatment
- Statistical analyses included descriptive measures such as mean ± standard deviation (SD), frequency, and percentage distributions
- HAE-related healthcare costs associated with outpatient visits, inpatient admissions, emergency room visits (ER), and home healthcare visits were described per patient per year (PPPY) with 95% confidence intervals (CI) for patients with ≥1 claim in each domain

Figure 1. Analytic Cohort Selection of Commercially-Insured Patients Receiving Non-Androgen LTPs



References

- Busse PJ, Christiansen SC, Riedl MA, et al. *J Allergy Clin Immunol Pract.* 2021;9(1):132-150.e3. doi:10.1016/j.jaip.2020.08.046
- Tachdjian R, et al. *Allergy Asthma Proc.* 2020;41(3):172-182


Results

Table 1. Patient Characteristics (Analytic Cohort)

	Total (N=210)	Lanadelumab (n=106)	SC C1-INH (n=71)	IV C1-INH (n=17)	Berotralstat (n=16)
Age on Index Date (years)					
Mean ± SD	40.9 ± 14.1	42.1 ± 14.4	40.1 ± 13.4	35.5 ± 13.1	43.0 ± 15.5
Median (IQR)	41.7 (30.1, 52.1)	43.8 (30.2, 52.4)	39.7 (30.8, 51.1)	35.1 (26.0, 44.3)	49.9 (30.3, 53.1)
Female, n (%)	152 (72.4)	76 (71.7)	50 (70.4)	16 (94.1)	10 (62.5)
Geographic Region, n (%)					
South	100 (47.6)	54 (50.9)	33 (46.5)	8 (47.1)	5 (31.3)
Midwest	43 (20.5)	22 (20.8)	12 (16.9)	5 (29.4)	4 (25.0)
West	44 (21.0)	17 (16.0)	20 (28.2)	1 (5.9)	6 (37.5)
Northeast	22 (10.5)	12 (11.3)	6 (8.5)	3 (17.6)	1 (6.3)
Unknown/missing	1 (0.5)	1 (0.9)	0 (0.0)	0 (0.0)	0 (0.0)
Year of Index, n (%)					
2017*	10 (4.8)	0 (0.0)	3 (4.2)	7 (41.2)	0 (0.0)
2018	59 (28.1)	20 (18.9)	32 (45.1)	7 (41.2)	0 (0.0)
2019	55 (26.2)	42 (39.6)	12 (16.9)	1 (5.9)	0 (0.0)
2020	26 (12.4)	15 (14.2)	10 (14.1)	0 (0.0)	1 (6.3)
2021	60 (28.6)	29 (27.4)	14 (19.7)	2 (11.8)	15 (93.8)
Top 5 Comorbidities, n (%)					
Gastrointestinal disorders	68 (32.4)	39 (36.8)	19 (26.8)	5 (29.4)	5 (31.3)
Anxiety or depression	53 (25.2)	32 (30.2)	12 (16.9)	6 (35.3)	3 (18.8)
Hypertension	40 (19.0)	19 (17.9)	12 (16.9)	4 (23.5)	5 (31.3)
Cardiovascular diseases	37 (17.6)	21 (19.8)	11 (15.5)	2 (11.8)	3 (18.8)
Autoimmune diseases	32 (15.2)	14 (13.2)	10 (14.1)	3 (17.6)	5 (31.3)

C1-INH, C1 inhibitor; ER, emergency room; HAE, hereditary angioedema; HRU, healthcare resource utilization; IV, intravenous; PPPY, per patient year; SC, subcutaneous.

Figure 2. Rate of HAE-Related HRU PPPY, Among Patients with ≥1 Visit



C1-INH, C1 inhibitor; ER, emergency room; HAE, hereditary angioedema; HRU, healthcare resource utilization; IV, intravenous; PPPY, per patient year; SC, subcutaneous.

- During a median 16-month follow-up, nearly all patients (95%) had ≥1 all-cause outpatient visit, and most (90%) had HAE-related outpatient visits; allergists were the most commonly seen specialists (69%)
- Approximately one-third of patients had ≥1 HAE-related ER visits during follow-up (33%), and nearly one-quarter (22%) had ≥1 HAE-related home health visit
- The highest rate of HAE-related HRU was observed for home health visits (among patients with ≥1 visit: 5.1 PPPY, 95% CI 4.7, 5.6) (Figure 2)

Table 2. HAE-Related Healthcare Costs Among Commercially-Insured Patients Treated with Non-Androgen LTP

	Total (N=210)	Lanadelumab (n=106)	SC C1-INH (n=71)	IV C1-INH (n=17)	Berotralstat (n=16)
Total Healthcare Costs, Mean Costs PPPY (95% CI)	\$641,166 (\$561,682, \$736,853)	\$588,755 (\$518,477, \$676,818)	\$718,374 (\$527,344, \$955,253)	\$698,343 (\$420,001, \$1,055,464)	\$585,030 (\$466,017, \$708,281)
Pharmacy Costs, Mean Costs PPPY (95% CI)	\$579,585 (\$508,662, \$664,653)	\$573,719 (\$504,731, \$662,329)	\$630,592 (\$451,076, \$860,339)	\$415,580 (\$222,327, \$642,922)	\$566,354 (\$447,902, \$692,069)
Outpatient Costs, Patients with ≥1 Visit, Mean Costs PPPY (95% CI)	\$3,297 (\$1,263, \$6,914)	\$3,995 (\$749, \$11,600)	\$1,371 (\$800, \$2,059)	\$7,178 (\$1,614, \$15,424)	\$3,312 (\$421, \$7,317)
Inpatient Costs, Patients with ≥1 Visit, Mean Costs PPPY (95% CI)	\$30,061 (\$11,708, \$55,763)	\$18,140 (\$3,222, \$38,836)	\$41,127 (\$10,614, \$92,733)	\$3,873 (\$3,873, \$3,873)	\$42,673 (\$42,673, \$42,673)
Emergency Room Costs, Patients with ≥1 Visit, Mean Costs PPPY (95% CI)	\$25,606 (\$4,085, \$56,763)	\$2,394 (\$1,310, \$3,729)	\$67,947 (\$6,637, \$158,046)	\$4,077 (\$820, \$8,360)	\$4,517 (\$1,403, \$8,323)
Home Health Costs, Patients with ≥1 Visit, Mean Costs PPPY (95% CI)	\$207,784 (\$92,170, \$366,674)	\$53,253 (\$12,130, \$103,232)	\$220,455 (\$38,325, \$535,451)	\$581,046 (\$172,967, \$1,088,598)	\$63,583 (\$56, \$188,171)

- Total HAE-related healthcare care costs (\$718,374), pharmacy costs (\$630,592), and emergency room costs (\$67,947) were highest for patients treated with SC C1-INH (Table 2)
- Patients using IV C1-INH experienced the highest outpatient costs (\$7,178) and home health costs (\$581,046) in patients that had ≥1 outpatient or home visit, respectively
- Inpatient costs were highest for the berotralstat cohort (\$42,673) in patients that had ≥1 inpatient visit

Conclusions

- This large retrospective insurance claims database study revealed that HAE-related resource utilization and costs were substantial, despite the use of LTP treatments
- Most (90%) patients had HAE-related outpatient visits and approximately one-third of patients had ≥1 HAE-related emergency room visit
- Home health visits were the most frequently used HAE-related HRU for patients with ≥1 visit
- Total HAE-related healthcare care costs, pharmacy costs, and emergency room costs were highest for patients treated with SC C1-INH

Acknowledgments

The authors wish to thank Jason Allaire, PhD, of Generativity Health Outcomes Research for his assistance with this poster. Funding for Dr. Allaire was provided by KalVista Pharmaceuticals.

Presented

Eastern Allergy Conference 2024, May 30 – June 2, in Palm Beach, FL

To view this poster after the presentation, visit KalVista Virtual Medical Booth

