

Barriers to Timely On-Demand Treatment of Hereditary Angioedema Attacks in Italian Patients

Mauro Cancian¹, Paola Triggianese², Pietro Accardo³, Francesco Arcoleo³, Donatella Bignardi⁴, Caterina Colangelo⁵, Francesco Giardino⁶, Antonio Gidaro⁷, Marica Giliberti⁸, Maria Domenica Guarino⁹, Paola Lucia Minciullo¹⁰, Stefania Nicola¹¹, Francesca Perego¹², Riccardo Senter¹³, Giuseppe Spadaro¹⁴, Massimo Triggiani¹⁵, Sherry Danese¹⁶, Julie Ulloa¹⁶, Vibha Desai¹⁷, Paul Audhya¹⁷, Andrea Zanichelli^{18,19}

¹Azienda Ospedale Università di Padova, Padova, Italy; ²Policlinico Universitario Tor Vergata, Department of Biomedicine and Prevention, Rome, Italy; ³A.O. "Ospedali Riuniti Villa Sofia-Cervello" – Presidio Ospedaliero Cervello, Palermo, Italy; ⁴IRCCS Ospedale Policlinico San Martino Genova, Genova, Italy; ⁵Azienda Sanitaria Locale di Pescara, Pescara, Italy; ⁶A.O.U. Policlinico "G. Rodolico-San Marco," Catania, Italy; ⁷Ospedale Luigi Sacco, Università degli Studi di Milano, Milano, Italy; ⁸Azienda Ospedaliero-Universitaria "Policlinico" di Bari, Bari, Italy; ⁹Presidio Ospedaliero di Civitanova Marche, Civitanova Marche, Italy; ¹⁰OU Policlinico "G. Martino" di Messina, Messina, Italy; ¹¹Allergy and Immunology Unit - AO Ordine Mauriziano di Torino and Department of Medical Sciences - University of Turin, Italy; ¹²IRCCS Istituti Clinici Scientifici Maugeri, Milano, Italy; ¹³Azienda Ospedaliera, Università degli Studi di Padova, Padova, Italy; ¹⁴Azienda Ospedaliera Universitaria Federico II di Napoli, Napoli, Italy; ¹⁵Azienda Ospedaliera Universitaria, Salerno, Italy; ¹⁶Outcomes Insights, Agoura Hills, CA, United States; ¹⁷KalVista Pharmaceuticals, Cambridge, MA, United States; ¹⁸Operative Unit of Medicine, Angioedema Center, IRCCS Policlinico San Donato, San Donato Milanese, Milan, Italy; ¹⁹Department of Biomedical Sciences for Health, University of Milan, Milan, Italy

Introduction

- Hereditary angioedema (HAE) is characterized by unpredictable swelling attacks affecting mucosal and subcutaneous tissues, which are typically painful, debilitating, and potentially fatal
- WAO/EAACI guidelines recommend the early use of on-demand treatment following recognition of an HAE attack to reduce morbidity and prevent mortality¹⁻³
- Despite the recommendation for early treatment, recent research suggests that patients delay on-demand treatment of their attacks⁴

Methods

- Individuals with Type 1 or 2 HAE due to C1 inhibitor deficiency were recruited through the Italian Network for Hereditary and Acquired Angioedema (ITACA) between September 2023 and January 2024
- Respondents enrolled were ≥12 years old and had to have treated with an approved on-demand therapy ≥1 HAE attack within 3 months prior to the survey
- The survey was self-reported, and took respondents approximately 20 minutes to complete
- Recruitment was stratified to include 50% of participants taking on-demand treatment only and 50% taking on-demand treatment + long-term prophylaxis (LTP)

Results

Table 1. Respondent Characteristics

	Total (n = 101)	On-Demand Only (n = 48)	On-Demand + LTP (n = 53)	Adults (n = 87)	Adolescents (n = 14)
Current Mean Age, Years (SD)	38 (16.2)	40 (16.5)	37 (15.9)	42 (14.0)	15 (1.6)
Diagnosis Mean Age, Years (SD)	17 (14.7)	20 (16.8)	14 (11.7)	18 (15.2)	7 (3.1)
Gender					
Male	39.6%	43.8%	35.8%	37.9%	50.0%
Female	60.04%	56.2%	64.2%	62.1%	50.0%
HAE Type					
Type 1	93.1%	93.8%	92.4%	93.1%	92.8%
Type 2	6.9%	6.2%	7.6%	6.9%	7.2%
Days Since Last Attack, Mean (SD)	19.0 (17.7)	17.6 (16.5)	20.2 (18.9)	18.3 (15.6)	23.0 (28.1)

LTP: long-term prophylaxis

- Respondents included 14 adolescents (14%) with an average age of 15 years and 87 adults (86%) with an average age of 42 years (Table 1)
- Overall, respondents were predominately female (60.04%) with an average of 19 days since last HAE attack

Acknowledgments

The authors wish to thank Jason Allaire, PhD of Generativity Health Outcomes Research for his assistance with this poster. Funding for Dr. Allaire was provided by KalVista Pharmaceuticals.

Disclosures

Gidaro Antonio was a speaker for Takeda and CSL Behring. Francesco Arcoleo received consultancy fees from Takeda, CSL Behring, BioCryst, and participated in clinical trials with Takeda, BioCryst, Ionis, Kalvista, Pharvaris. Paul Audhya is an employee of and owns stock in Kalvista. Mauro Cancian received honoraria and/or meeting/travel support paid to the institution from KalVista Pharmaceuticals, BioCryst, CSL Behring, Pharvaris, and Takeda. Sherry Danese received consulting fees from Kalvista. Vibha Desai is an employee of and owns stock in Kalvista. Francesco Giardino served on advisory boards/seminars funded by BioCryst, CSL Behring, Kalvista, Takeda and received funding to attend conferences/educational events from CSL Behring, Takeda, Marica Giliberti is a consultant for Takeda, Sanofi Genzyme, Chiesi, AstraZeneca, BioCryst, CSL Behring, Kyowa Kirin, Alnylam. Francesca Perego participated in clinical trials for Takeda. Advisory boards for BioCryst, Takeda, and CSL Behring. Riccardo Senter served as a consultant for BioCryst and Takeda and received travel grants from Takeda, BioCryst, CSL Behring, Aik Abello, Novartis. Massimo Triggiani received fees for advisory board from Takeda, advisory board fees for CSL Behring and BioCryst. Julie Ulloa received consulting fees from KalVista. Andrea Zanichelli received honoraria, meeting/travel support, and/or served on advisory boards for KalVista Pharmaceuticals, Astria, BioCryst, CSL Behring, Pharming, Pharvaris, and Takeda.

Results

Figure 1. On-Demand Treatment at Time of Last Treated Attack

Treatment Used (n=101)	On-Demand Only (n=48)	On-Demand + LTP (n=53)	Adults (n=87)	Adolescents (n=14)
Icatibant (Firazyr and generic)	55%	55%	59%	29%
Plasma derived C1 esterase inhibitor (Berinert)	44%	42%	39%	71%
Plasma derived C1 esterase inhibitor (Cinryze)	2%	4%	2%	–
Recombinant C1 esterase inhibitor (Ruconest)	0%	–	–	–


- The most commonly used initial on-demand treatment was icatibant (branded and generic) for adults and plasma derived C1 esterase inhibitor (Berinert) for adolescents (Figure 1)
- Among both the on-demand only and on-demand plus long-term prophylaxis groups, icatibant (branded and generic) was the most frequently used treatment, closely followed by plasma derived C1 esterase inhibitor (Berinert)

Figure 2. Long-Term Prophylaxis at Time of Last Treated Attack

	Adults (n = 44)	Adolescents (n = 9)
Plasma derived C1 esterase inhibitor (Berinert)	34%	67%
Lanadelumab	30%	11%
Berotrastat	11%	11%
Danazol	11%	–
Plasma derived C1 esterase inhibitor (Cinryze)	8%	11%
Tranexamic acid	6%	–


- Among those on long-term prophylaxis at the time of the last treated attack, plasma derived C1 esterase inhibitor (Berinert) was the most common treatment among adolescents, whereas adults were most often treated with lanadelumab (Figure 2)

Figure 3. Time to On-Demand Treatment After Attack Onset



- The mean time (SD) to treatment during the most recent attack was 2.9 hours (2.7), with 10% (10/101) treating in <1 hour (Figure 3)

Figure 4. Perception of Time to Treatment Versus Actual Time to Treatment for Those Who Perceived They Treated Early



- 71% of respondents (72/101) believed they treated their attack early, despite only 14% of them treating in less than one hour (Figure 4)
- The mean time to treatment for those who believed they treated early was 2.1 hours

Figure 5. Barriers to Treating Attack Early Excluding Those Who Treated the Attack Immediately (n = 81; excludes those who treated attacks immediately)

Barriers (Detailed) (Excluding those who treated the attack immediately, ranked top 5)	Ranked Top 5			
	On-Demand Only (n=40)	On-Demand + LTP (n=41)	Adults (n=70)	Adolescents (n=11)
I was not certain it was a real / actual attack	40%	38%	42%	39%
I thought the attack would be mild	37%	30%	44%	40%
I wanted to save my on-demand treatment for a severe attack	22%	18%	27%	23%
I waited to treat until the attack was severe	20%	18%	22%	19%
I did not want to / could not interrupt what I was doing	16%	23%	10%	16%
I did not have anyone to help me	12%	13%	12%	9%
I did not have my on-demand treatment with me	9%	10%	7%	7%
I did not have a private place to administer the treatment	9%	13%	5%	9%
I had to go to the hospital / emergency center for treatment	7%	13%	2%	7%
I wanted to avoid the burning, stinging or pain with injection	5%	5%	5%	3%
I wanted to avoid the pain of the needle	4%	3%	5%	3%
I wanted to avoid the side effects of treatment	3%	5%	–	3%
I did not feel well enough to prepare and administer the treatment	3%	3%	2%	3%
My on-demand treatment was expensive	1%	–	2%	1%

- Eighty-one respondents (80%) who did not treat immediately ranked their top 5 reasons for not treating earlier (Figure 5)
- The most common barriers to treating sooner were uncertainty the attack was real (40%), thinking the attack would remain mild (37%), and wanting to save on-demand treatment for a severe attack (22%)
- Treatment administration-related barriers (e.g., not wanting to interrupt what they were doing, not having anyone to help with administration) were reported by 38% of respondents as their top reason for delaying treatment

Conclusions

- Most respondents did not meet guideline recommendations for immediate on-demand treatment following HAE attack onset
- Uncertainty that the attack was real and thinking the attack was going to stay mild were the most common barriers to treating immediately
- Substantial proportion reported treatment administration barriers to treating the attack immediately, such as not wanting to interrupt what they were doing, not having anyone to help, and not having a private place to administer treatment
- These findings highlight a need to proactively address barriers contributing to treatment delays, including a need for oral on-demand treatment option

Presented

AAAAI Annual Meeting 2025.
February 28 to March 3, 2025 in San Diego, CA.

Please scan the QR code to view this poster after the presentation.



References

- Betschel S, Badiou J, Binkley K, et al. The International/Canadian Hereditary Angioedema Guideline. *Allergy, Asthma & Clinical Immunology*. 2019/11/25 2019;15(1):72. doi:10.1186/s13223-019-0376-8
- Busse PJ, Christiansen SC, Riedl MA, et al. US HAEA Medical Advisory Board 2020 Guidelines for the Management of Hereditary Angioedema. *The Journal of allergy and clinical immunology in practice*. Jan 2021;9(1):132-150.e3. doi:10.1016/j.jaip.2020.08.046
- Maurer M, Mager M, Betschel S, et al. The international WAO/EAACI guideline for the management of hereditary angioedema-The 2021 revision and update. *Allergy*. Jul 2022;77(7):1951-1990. doi:10.1111/all.15214
- Radjoicic, Cristine et al. Patient Perspectives On Early Use Of On-demand Treatment For Hereditary Angioedema (HAE) Attacks to Reduce Severity and Duration. *Journal of Allergy and Clinical Immunology*, (2023) Volume 151, Issue 2, AB143